

Poisson distribution

A stochastic experiment is repeated many times. Let the expected number of successes be ${\n}^{{\n}}_{{\n}}$. Then the probability of observing n successes would be

$$p_{\ln(n)={{\ln^ne^{-\ln }}}$$

The Poisson distribution is of interest especially for TCSPC: The expected number of photons in any TCSPC channel is given by the 'real' decay (including convolution with the IRF etc.), while the stochastic nature of the measurement process (either a photon is detected or it is not) introduces noise, which follows a Poisson distribution. In the limit of large $\{\nu\}^{{}_{1}}$ the Poisson distribution approaches a WGaussian distribution with a width of ν centred around $\{\nu\}^{{}_{1}}$.

In the Gaussian limit least squares Wfitting may be applied, otherwise MLE Wfitting is preferable.

Copyright of this document belongs to PicoQuant GmbH. No parts of it may be reproduced, translated or transferred to third parties without written permission of PicoQuant GmbH. All information given here is reliable to our best knowledge. However, no responsibility is assumed for possible inaccuraciesor omissions. Specifi cations and external appearances are subject to change without notice.



PicoQuant GmbH Rudower Chaussee 29 (IGZ) 12489 Berlin Germany P +49-(0)30-1208820-89 F +49-(0)30-1208820-90 info@picoquant.com www.picoquant.com

© PicoQuant GmbH 1/1