

## Residuals

In the context of [least squares](#) (or any other [fitting](#) method, e.g. [MLE](#)) the residuals are the difference between a model function and the experimental data. In [least squares](#) usually the weighted residuals trace is shown:

$$R_{\text{wt}}(t_i) = \frac{D(\text{model parameters}, t_i) - D_i^{\text{exp}}}{w_i}$$

$D_i^{\text{exp}}$  is the  $i$ -th data point of an experimental data set,  $D(\text{model parameters}, t_i)$  is the model equation at the observed points  $t_i$ .

The [weighting](#) factor  $w_i$  describes the experimental uncertainty of each individual data point. For [TCSPC](#) data  $w_i$  is defined as

$$w_i = \sqrt{D_i^{\text{exp}}}$$

The residuals trace is of importance within any framework concerned with fitting, as the [SymPhoTime](#) software or [FluoFit](#).

Copyright of this document belongs to PicoQuant GmbH. No parts of it may be reproduced, translated or transferred to third parties without written permission of PicoQuant GmbH. All information given here is reliable to our best knowledge. However, no responsibility is assumed for possible inaccuracies or omissions. Specifications and external appearances are subject to change without notice.



PicoQuant GmbH  
Rudower Chaussee 29 (IGZ)  
12489 Berlin  
Germany

P +49-(0)30-1208820-89  
F +49-(0)30-1208820-90  
info@picoquant.com  
www.picoquant.com